

A CANONICAL DECOMPOSITION FOR LINEAR OPERATORS AND LINEAR RELATIONS

S. HASSI, Z. SEBESTYÉN, H.S.V. DE SNOO, AND F.H. SZAFRANIEC

ABSTRACT. An arbitrary linear relation (multivalued operator) acting from one Hilbert space to another Hilbert space is shown to be the sum of a closable operator and a singular relation whose closure is the Cartesian product of closed subspaces. This decomposition can be seen as an analog of the Lebesgue decomposition of a measure into a regular part and a singular part. The two parts of a relation are characterized metrically and in terms of Stone's characteristic projection onto the closure of the linear relation.

1. INTRODUCTION

Let T be a linear operator from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. In general the closure $\overline{T}$ of the operator T (i.e., the closure of the graph of T in the Cartesian product $\mathfrak{H} \times \mathfrak{K}$) is not the graph of an operator anymore. In other words, (the graph of) $\overline{T}$ has a nontrivial multivalued part $\text{mul } \overline{T} = \{k \in \mathfrak{K} : \{0, k\} \in \overline{T}\}$. Relative to the closed linear subspace $\text{mul } \overline{T}$ of $\mathfrak{H}$, P.E.T. Jorgensen [11] and S. Ôta [20], [21] have given a decomposition of a densely defined operator T as an operator sum of a *closable* operator, whose closure is again (the graph of) an operator, and a singular operator, whose closure is the Cartesian product of a closed subspace of $\mathfrak{H}$ and a closed subspace of $\mathfrak{K}$. This decomposition is similar to a decomposition of nonnegative bounded linear operators due to T. Ando [1] (see also [13], [18]) and a decomposition of semibounded sesquilinear forms due to B. Simon (see [14], [25], [26]). It was pointed out in these publications that there is an analogy with the Lebesgue decomposition of a measure into a regular part and a singular part.

The purpose of this note is to show that there is a similar decomposition in the case of linear relations from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. The notion of a linear relation as a multivalued linear operator was introduced by R. Arens [3] and extensively studied by E.A. Coddington [4] and by many others. The treatment of Jorgensen and Ôta for operators can be relaxed: it is not necessary to consider operators which are densely defined and in fact their treatment remains true for relations. Indeed the language of relations seems to be the proper context for such decompositions. Now the result is that any linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ has a decomposition as an operator-like sum of a closable operator whose closure is again (the graph of) an operator and a singular relation whose closure is the Cartesian product of a closed subspace of $\mathfrak{H}$ and a closed subspace of $\mathfrak{K}$. The components of this decomposition can be characterized in various ways.

2000 *Mathematics Subject Classification.* Primary 47A05, 47A06.

Key words and phrases. Relation, multivalued operator, graph, adjoint relation, closable operator, regular relation, singular relation, Stone decomposition.

The research was supported by bilateral agreements between the Eötvös Loránd University in Budapest, the University of Groningen, and the Jagiellonian University in Krakow, and by the Research Institute for Technology of the University of Vaasa. The fourth author was also supported by the KBN grant 2 PO3A 037 024.

When the relation T from $\mathfrak{H}$ to $\mathfrak{K}$ itself is considered with the graph inner product, then its completion can be contractively embedded into the Hilbert space $\mathfrak{H}$; the kernel of this contraction corresponds to the multivalued part $\text{mul } \overline{T}$. This observation leads to a metric characterization of the decomposition of the relation T . There is a similar description for the decomposition of a pair of nonnegative sesquilinear forms, cf. [9]. By means of the above mentioned result a metric characterization of closable operators is presented.

The decomposition of Jorgensen and Ôta also has connections with the characteristic (projection) matrix introduced by J. von Neumann [17] and M.H. Stone [28], and Stone's decomposition of a linear operator [28]. The work of A.E. Nussbaum [19] concerning orthogonal projections onto closed subspaces of a Cartesian product is easily translated for closed linear relations and this leads to the Stone decomposition for closed linear relations, cf. [16]. This makes it possible to characterize the regular and singular parts of a linear relation T from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ in terms of the orthogonal projection from the Cartesian product $\mathfrak{H} \times \mathfrak{K}$ onto the closure $\overline{T}$.

It is also shown how the main decomposition result in this paper (see Theorem 4.1) can be obtained by applying the general characterization result of $\text{dom } T^*$ in [6] (cf. also Lemma 9.1 below); here T^* is the adjoint relation of the linear relation T .

2. PRELIMINARIES

Here is a short review of notions associated with linear relations. Recall that a linear relation T from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ is a (not necessarily closed) linear subspace of the Cartesian product $\mathfrak{H} \times \mathfrak{K}$. The domain, range, kernel, and multivalued part of a linear relation T are defined by:

$$\begin{aligned} \text{dom } T &\stackrel{\text{def}}{=} \{f \in \mathfrak{H} : \{f, f'\} \in T\}, \quad \text{ran } T \stackrel{\text{def}}{=} \{f' \in \mathfrak{K} : \{f, f'\} \in T\}, \\ \ker T &\stackrel{\text{def}}{=} \{f \in \mathfrak{H} : \{f, 0\} \in T\}, \quad \text{mul } T \stackrel{\text{def}}{=} \{f' \in \mathfrak{K} : \{0, f'\} \in T\}. \end{aligned}$$

The formal inverse T^{-1} is a linear relation from $\mathfrak{K}$ to $\mathfrak{H}$ which is obtained from T by interchanging the components of the elements of T . Clearly, $\text{ran } T = \text{dom } T^{-1}$ and $\text{mul } T = \ker T^{-1}$. The linear relation T is said to be closed if T is a closed subspace of the Cartesian product $\mathfrak{H} \times \mathfrak{K}$. If T is closed the kernel $\ker T$ and multivalued part $\text{mul } T$ of T are automatically closed. The closures of the domain and range of a linear relation T are denoted by $\overline{\text{dom } T}$ and $\overline{\text{ran } T}$. Observe that

$$(2.1) \quad \text{dom } \overline{T} \subset \overline{\text{dom } T} \quad \text{and} \quad \text{ran } \overline{T} \subset \overline{\text{ran } T}.$$

To see this, let $g \in \text{ran } \overline{T}$, so that $\{f, g\} \in \overline{T}$ for some $f \in \mathfrak{H}$. Then there exist elements $\{f_n, g_n\} \in T$ such that $\{f_n, g_n\} \rightarrow \{f, g\}$. This shows that $g \in \overline{\text{ran } T}$ and the second inclusion (2.1) follows. The first identity is obtained from the second one by inverting the relation T . The adjoint T^* is a closed linear relation from $\mathfrak{K}$ to $\mathfrak{H}$ defined by $T^* = JT^\perp = (JT)^\perp$, where $J\{f, f'\} = \{f', -f\}$. Observe that $T^{**} = (T^\perp)^\perp$ is the closure $\overline{T}$ of T in $\mathfrak{H} \times \mathfrak{K}$. Each of the following identities is clear:

$$(2.2) \quad (\text{ran } T)^\perp = \ker T^*, \quad (\text{dom } T)^\perp = \text{mul } T^*.$$

Furthermore, the one identity in (2.2) is obtained by inverting the relation in the other identity in (2.2). When T in the identities in (2.2) is replaced by its adjoint T^* , one also obtains

$$(2.3) \quad (\text{ran } T^*)^\perp = \ker \overline{T}, \quad (\text{dom } T^*)^\perp = \text{mul } \overline{T}.$$

For two linear relations A and B from $\mathfrak{H}$ to $\mathfrak{K}$ there is a *componentwise sum* $A \hat{+} B$ from $\mathfrak{H}$ to $\mathfrak{K}$, defined by

$$A \hat{+} B \stackrel{\text{def}}{=} \{\{f + g, f' + g'\} : \{f, f'\} \in A, \{g, g'\} \in B\}.$$

Note that $^1 (A \hat{+} B)^* = A^* \cap B^*$ and $(A \hat{+} B)^- = (\overline{A} \hat{+} \overline{B})^-$. The notation $A \hat{+} B$ is used to indicate that A and B are orthogonal in the Cartesian product. As an example, observe that if A is a linear relation from $\mathfrak{H}$ to $\mathfrak{K}$ and $\mathfrak{R}$ is a linear subspace of $\mathfrak{K}$, then the relation T from $\mathfrak{H}$ to $\mathfrak{K}$, defined by

$$(2.4) \quad T = \{ \{f, f' + \varphi\} : \{f, f'\} \in A, \varphi \in \mathfrak{R} \},$$

can be written as a componentwise sum $T = A \hat{+} B$, where the linear relation B from $\mathfrak{H}$ to $\mathfrak{K}$ is defined by $B = \{ \{0, \varphi\} : \varphi \in \mathfrak{R} \}$. This leads to

$$(2.5) \quad \overline{T} = (\overline{A} \hat{+} \{ \{0, \varphi\} : \varphi \in \mathfrak{R}^- \})^-.$$

There is also an *operator-like sum* $A + B$ from $\mathfrak{H}$ to $\mathfrak{K}$, defined by

$$A + B \stackrel{\text{def}}{=} \{ \{f, f' + f''\} : \{f, f'\} \in A, \{f, f''\} \in B \},$$

so that $\text{dom}(A + B) = \text{dom } A \cap \text{dom } B$. The operator-like sum reduces to the usual operator sum if A and B are (graphs of) operators.

Now let A be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{R}$, and let B be a linear relation from the Hilbert space $\mathfrak{R}$ to a Hilbert space $\mathfrak{K}$. The product BA is a linear relation from $\mathfrak{H}$ to $\mathfrak{K}$, defined by

$$BA \stackrel{\text{def}}{=} \{ \{f, f'\} : \{f, g\} \in A, \{g, f'\} \in B \}.$$

In general, $A^*B^* \subset (BA)^*$. However, when B or A^{-1} is (the graph of) a bounded everywhere defined operator on $\mathfrak{R}$, then

$$(2.6) \quad (BA)^* = A^*B^*.$$

This is known for the case of operators; for a proof in the case of linear relations, see for instance [8].

3. REGULAR AND SINGULAR RELATIONS

A classical result of J. von Neumann states that the adjoint of a densely defined linear operator is densely defined if and only if the operator itself is closable (i.e., its closure is an operator). The result which follows, and which can be deduced immediately from the second identity in (2.3), does not require the object in question to be *a priori* (the graph of) an operator; the latter comes out as an additional conclusion. Thus call a linear relation T from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ *regular* if its closure $\overline{T}$ is (the graph of) an operator.

Proposition 3.1. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then the following statements are equivalent:*

- (i) $\overline{T}$ is regular;
- (ii) $\text{dom } T^* = \mathfrak{K}$,

in which case T is (the graph of) an operator.

Note that a linear operator T is regular if and only if it is closable. If a linear relation T is regular, then T is automatically a closable operator. For any linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ the adjoint relation T^* from $\mathfrak{K}$ to $\mathfrak{H}$ is a closed linear relation; moreover, the adjoint T^* is a closed operator if and only if $\text{dom } T$ is dense in $\mathfrak{H}$ (independent of T being regular).

A linear relation T from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ is said to be *singular* if

$$(3.1) \quad \text{ran } T \subset \text{mul } \overline{T} \quad \text{or equivalently} \quad \overline{\text{ran } T} \subset \text{mul } \overline{T}.$$

The equivalence here is due to the closedness of $\text{mul } \overline{T}$. Furthermore, the inclusion

$$(3.2) \quad \text{mul } \overline{T} \subset \overline{\text{ran } T},$$

¹ A dash $-$ put aside still refers to the closure operation.

follows from (2.1) as $\text{mul } \overline{T} \subset \text{ran } \overline{T}$. Therefore, a linear relation T is singular if and only if

$$(3.3) \quad \overline{\text{ran}} T = \text{mul } \overline{T},$$

which follows by combining (3.1) and (3.2).

Proposition 3.2. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then the following statements are equivalent:*

- (i) T is singular;
- (ii) $\text{dom } T^* \subset \ker T^*$ or, equivalently, $\text{dom } T^* = \ker T^*$;
- (iii) $T^* = \text{dom } T^* \times \text{mul } T^*$;
- (iv) $\overline{T} = \overline{\text{dom } T} \times \text{mul } \overline{T}$.

Proof. (i) $\Rightarrow$ (ii) The identity in (3.3) implies that $(\overline{\text{ran}} T)^\perp = (\text{mul } \overline{T})^\perp$, which is equivalent to $\ker T^* = \overline{\text{dom } T^*}$ by (2.2) and (2.3). This implies that $\text{dom } T^* \subset \ker T^*$, while the reverse inclusion is obvious.

(ii) $\Rightarrow$ (iii) Let $\{f, g\} \in T^*$. Then in particular $f \in \text{dom } T^*$ and so $f \in \ker T^*$ by (ii). Therefore $\{f, 0\} \in T^*$ and this implies that $\{0, g\} \in T^*$, or $g \in \text{mul } T^*$. This shows that $\{f, g\} \in \text{dom } T^* \times \text{mul } T^*$. Conversely, let $\{f, g\} \in \text{dom } T^* \times \text{mul } T^*$. Then $\{0, g\} \in T^*$ by definition. Moreover, $f \in \text{dom } T^*$ or by (ii), $f \in \ker T^*$, so that $\{f, 0\} \in T^*$. Clearly, by linearity, this implies that $\{f, g\} \in T^*$.

(iii) $\Rightarrow$ (iv) Taking adjoints in (iii) yields $T^{**} = (\text{mul } T^*)^\perp \times (\text{dom } T^*)^\perp$, which gives (iv) by means of (2.2) and (2.3).

(iv) $\Rightarrow$ (i) It follows from $\text{ran } \overline{T} = \text{mul } \overline{T}$ that $\text{ran } T \subset \text{mul } \overline{T}$. Hence, by definition, T is singular. $\square$

Corollary 3.3. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then the following statements are equivalent:*

- (i) T is singular;
- (ii) T^{-1} is singular;
- (iii) T^* is singular.

Proof. (i) $\Leftrightarrow$ (ii) Assume that T is singular. Then part (iv) of Proposition 3.2 implies that

$$(3.4) \quad \ker \overline{T} = \overline{\text{dom } T} \quad \text{or equivalently} \quad \text{mul } \overline{T^{-1}} = \overline{\text{ran}} T^{-1}.$$

Hence, T^{-1} is singular by (3.3). For the reverse implication it is now enough to observe that $(T^{-1})^{-1} = T$.

(ii) $\Leftrightarrow$ (iii) Part (ii) of Proposition 3.2 shows that T^* is singular if and only if $\text{dom } \overline{T} = \ker \overline{T}$. Clearly this is equivalent to (3.4), which means that T^{-1} is singular. $\square$

Note that if T is singular, then in particular $\text{dom } T^* = \ker T^*$ is closed. A linear relation T from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ is said to be *maximally singular* if

$$\text{dom } T^* = \{0\} \quad \text{or equivalently} \quad \text{mul } \overline{T} = \mathfrak{K}.$$

Proposition 3.4. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then the following statements are equivalent:*

- (i) T is maximally singular;
- (ii) T is singular and $\overline{\text{ran}} T = \mathfrak{K}$;
- (iii) $\overline{T} = \overline{\text{dom } T} \times \mathfrak{K}$.

Proof. (i) $\Rightarrow$ (ii) Assume that T is maximally singular. Clearly

$$\text{ran } T \subset \mathfrak{K} = \text{mul } \overline{T} \subset \text{ran } \overline{T} \subset \overline{\text{ran}} T,$$

where the last inclusion follows from (2.1). Hence (3.1) is satisfied and $\overline{\text{ran}} T = \mathfrak{K}$.

(ii) $\Rightarrow$ (iii) Apply part (iv) of Proposition 3.2 with $\text{mul } \overline{T} = \overline{\text{ran}} T = \mathfrak{K}$.

(iii) $\Rightarrow$ (i) Clearly, $\text{mul } \overline{T} = \mathfrak{K}$, so that T is maximally singular. $\square$

As a simple example, observe that the linear relation T defined by $T = K^*U$ is maximally singular, when U is any linear relation and K is an injective bounded linear operator with $\text{ran } K \subset (\text{dom } U^*)^\perp$. To see this, note that $T^* = U^*K$, cf. (2.6). Hence, if $\{f, f'\} \in T^*$, then $\{f, Kf\} \in K$ and $\{Kf, f'\} \in U^*$. This shows that $Kf \in \text{dom } U^*$ which leads to $Kf = 0$. Since K is injective, it follows that $f = 0$. Therefore $\text{dom } T^* = \{0\}$ and T is maximally singular, cf. [21].

Observe that V.D. Koshmanenko and S. Ôta [15] consider (densely defined) linear operators T which satisfy the property $\text{dom } T \subset \ker \overline{T}$. The inverse T^{-1} of such an operator is singular in the present sense, see (3.1), and according to Corollary 3.3 then equivalently T is singular.

4. CANONICAL DECOMPOSITIONS

Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ and let $\overline{T}$ be its closure in the Cartesian product $\mathfrak{H} \times \mathfrak{K}$. Denote the orthogonal projection from $\mathfrak{K}$ onto $\text{mul } \overline{T}$ by P . With T are associated the linear relation T_{reg} from $\mathfrak{H}$ to $\mathfrak{K}$ defined by

$$(4.1) \quad T_{\text{reg}} \stackrel{\text{def}}{=} \{ \{f, (I - P)f'\} : \{f, f'\} \in T \},$$

and the linear relation T_{sing} from $\mathfrak{H}$ to $\mathfrak{K}$ defined by

$$(4.2) \quad T_{\text{sing}} \stackrel{\text{def}}{=} \{ \{f, Pf'\} : \{f, f'\} \in T \}.$$

Observe that T_{reg} and T_{sing} have the same domain $\text{dom } T$. Moreover T_{reg} and T_{sing} are (graphs of) operators if T itself is (the graph of) an operator. The following decomposition result is an adaptation of a result of Jorgensen [11].

Theorem 4.1. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then T admits the canonical operator-like sum decomposition*

$$(4.3) \quad T = T_{\text{reg}} + T_{\text{sing}},$$

where T_{reg} is a regular relation from $\mathfrak{H}$ to $\mathfrak{K}$ and T_{sing} is a singular relation from $\mathfrak{H}$ to $\mathfrak{K}$ with

$$\text{mul } T_{\text{sing}} = \text{mul } T, \quad \text{mul } (T_{\text{sing}})^- = \text{mul } \overline{T}.$$

Proof. To show that the relation T_{reg} in (4.1) is regular, it suffices to show that its closure is an operator. Assume therefore that there is a sequence $\{f_n, f'_n\} \in T$ such that

$$\{f_n, (I - P)f'_n\} \mapsto \{0, g\},$$

which implies that $g \in (\text{mul } \overline{T})^\perp$. Furthermore, it follows from the definition of T^* that for all $\{h, h'\} \in T^*$

$$(4.4) \quad \begin{aligned} 0 &= (f'_n, h) - (f_n, h') \\ &= ((I - P)f'_n, h) - (f_n, h') + (Pf'_n, h) \\ &= ((I - P)f'_n, h) - (f_n, h'). \end{aligned}$$

Here the second identity in (2.3) has been used. Taking the limit $n \rightarrow \infty$ in (4.4) leads to $(g, h) = 0$ for all $h \in \text{dom } T^*$. Hence $g \in (\text{dom } T^*)^\perp = \text{mul } \overline{T}$. Therefore $g = 0$. It follows that T_{reg} is a closable operator.

Next it will be shown that the relation T_{sing} is singular. Note that $\{h, h'\} \in (T_{\text{sing}})^*$ if and only if

$$(h', f) = (h, Pf') = (Ph, f') \quad \text{for all } \{f, f'\} \in T,$$

or, equivalently, if and only if $\{Ph, h'\} \in T^*$. Therefore $h \in \text{dom } (T_{\text{sing}})^*$ if and only if $Ph \in \text{dom } T^*$. Now observe that $Ph \in \text{dom } T^*$ if and only if $Ph = 0$, since $\text{dom } T^* \subset (\text{mul } \overline{T})^\perp$. Furthermore, $Ph = 0$ is equivalent to $h \in (\text{mul } \overline{T})^\perp = \overline{\text{dom } T^*}$. Hence it follows that

$$(4.5) \quad \text{dom } (T_{\text{sing}})^* = \overline{\text{dom } T^*}.$$

The same argument shows that $h \in \ker (T_{\text{sing}})^*$ if and only if $Ph \in \ker T^*$. Now, if $Ph \in \ker T^*$ then $Ph \in \text{dom } T^*$ and $Ph = 0$. Conversely, if $Ph = 0$ then $Ph \in \ker T^*$. Hence, it follows that

$$(4.6) \quad \ker (T_{\text{sing}})^* = \overline{\text{dom } T^*}.$$

Combining (4.5) and (4.6) gives

$$\text{dom } (T_{\text{sing}})^* = \ker (T_{\text{sing}})^*,$$

in other words, the relation T_{sing} is singular by (ii) of Proposition (3.2).

Moreover, (4.5) shows that

$$\text{mul } (T_{\text{sing}})^- = (\text{dom } (T_{\text{sing}})^*)^\perp = (\overline{\text{dom } T^*})^\perp = \text{mul } \overline{T}.$$

Finally, observe that

$$\text{mul } T_{\text{sing}} = \{Pf' : \{0, f'\} \in T\} = \{f' : \{0, f'\} \in T\} = \text{mul } T.$$

This completes the proof. $\square$

Corollary 4.2. *The singular part of T_{reg} is the zero operator and the regular part of T_{sing} is the zero operator. Hence T_{reg} is equal to its regular part and T_{sing} is equal to its singular part. In particular, T is regular if and only if $T = T_{\text{reg}}$ and T is singular if and only if $T = T_{\text{sing}}$.*

Proof. The operator T_{reg} is closable; hence the multivalued part of its closure is trivial. Therefore, its singular part is the zero operator on $\text{dom } T$.

The multivalued part of the closure of T_{sing} is equal to $\text{mul } (T_{\text{sing}})^- = \text{mul } \overline{T}$. Hence, the regular part of T_{sing} is given by

$$\{f, (I - P)Pf'\} : \{f, f'\} \in T\},$$

which is the zero operator on $\text{dom } T$.

If $T = T_{\text{reg}}$ ($T = T_{\text{sing}}$), then T is regular (singular) by Theorem 4.1. Conversely, if T is regular then according to Proposition 3.1 $\overline{\text{dom } T^*} = \mathfrak{K}$ or, equivalently, $\text{mul } \overline{T} = \{0\}$. Thus, $P = 0$ and $T = T_{\text{reg}}$.

Finally, if T is singular then according to (3.3) $Pf' = f'$ for all $f' \in \overline{\text{ran } T}$. In particular, $Pf' = f'$ for all $f' \in \text{ran } T$ and therefore $T = T_{\text{sing}}$. $\square$

Corollary 4.3. *The singular part T_{sing} is maximally singular if and only if T is maximally singular.*

Proof. Observe that the identity $\text{mul } (T_{\text{sing}})^- = \text{mul } \overline{T}$ implies that T_{sing} is maximally singular if and only if T is maximally singular. $\square$

The canonical decomposition in Theorem 4.1 is about the decomposition of a linear relation T as an operator-like sum of a regular relation T_{reg} (a closable operator) and a singular relation T_{sing} . However, observe that because

$$\{f, f'\} = \{f, (I - P)f' + Pf'\}, \quad \{f, f'\} \in T,$$

it can also be written as a component-wise sum

$$\{f, f'\} = \{f, (I - P)f'\} \hat{+} \{0, Pf'\}, \quad \{f, f'\} \in T$$

which in fact is an orthogonal sum $\hat{+}$ in $\mathfrak{H} \times \mathfrak{K}$. This implies that

$$(4.7) \quad T \subset T_{\text{reg}} \hat{+} \{\{0, g\} : g \in \text{mul } \overline{T}\}.$$

When T is closed, this argument leads to the usual decomposition which goes back to [3] and [4].

Proposition 4.4. *Let T be a closed linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then $\text{mul } T$ is closed, T_{reg} is a closed operator, and*

$$(4.8) \quad T = T_{\text{reg}} \hat{\oplus} \{ \{0, g\} : g \in \text{mul } T \}.$$

Proof. If T is closed, then $\text{mul } \overline{T} = \text{mul } T$ is closed and P is an orthogonal projection from $\mathfrak{K}$ onto $\text{mul } T$. Hence in

$$\{f, f'\} = \{f, (I - P)f' + Pf'\}, \quad \{f, f'\} \in T,$$

one also has $\{0, Pf'\} \in T$, which leads to

$$\{f, (I - P)f'\} \in T \quad \text{for all} \quad \{f, f'\} \in T.$$

It follows that $T_{\text{reg}} \subset T$, and it is straightforward to see that

$$T_{\text{reg}} = \{ \{f, g\} \in T : g \perp \text{mul } T \},$$

which is clearly closed. Indeed, the righthand side of (4.8) is contained in the lefthand side. The reverse inclusion follows from (4.7) keeping in mind that $\text{mul } \overline{T} = \text{mul } T$. $\square$

For similar orthogonal operator parts under the weaker condition that only $\text{mul } T$ is closed, see [10].

Proposition 4.5. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then $(\overline{T})_{\text{reg}}$ is closed and*

$$(4.9) \quad (\overline{T})_{\text{reg}} = (T_{\text{reg}})^-.$$

Furthermore,

$$(4.10) \quad ((\overline{T})_{\text{sing}})^- = (T_{\text{sing}})^-.$$

Proof. The definition of regular part implies that $T_{\text{reg}} \subset (\overline{T})_{\text{reg}}$. To see this let $\{f, (I - P)f'\}$ with $\{f, f'\} \in T$ be an element in T_{reg} . But then also $\{f, f'\} \in \overline{T}$ and since P is an orthogonal projection onto $\text{mul } \overline{T}$, it follows that $\{f, (I - P)f'\}$ belongs to $(\overline{T})_{\text{reg}}$.

The definition of regular part also implies that $(\overline{T})_{\text{reg}} \subset (T_{\text{reg}})^-$. To see this let $\{f, (I - P)f'\}$ with $\{f, f'\} \in \overline{T}$ be an element of $(\overline{T})_{\text{reg}}$. Then there exists a sequence $\{f_n, f'_n\} \in T$ such that $\{f_n, f'_n\} \rightarrow \{f, f'\}$. However this implies that the sequence $\{f_n, (I - P)f'_n\} \in T_{\text{reg}}$ approximates $\{f, (I - P)f'\}$. In other words $(\overline{T})_{\text{reg}} \subset (T_{\text{reg}})^-$.

Combining these two assertions it follows that

$$T_{\text{reg}} \subset (\overline{T})_{\text{reg}} \subset (T_{\text{reg}})^-,$$

which leads to

$$((\overline{T})_{\text{reg}})^- = (T_{\text{reg}})^-.$$

A similar argument for the singular part gives

$$((\overline{T})_{\text{sing}})^- = (T_{\text{sing}})^-.$$

Therefore (4.10) follows. Moreover, since $\overline{T}$ is closed, it follows from Proposition 4.4 that $\overline{T}_{\text{reg}}$ is closed, which leads to (4.9). $\square$

Corollary 4.6. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then*

$$(4.11) \quad (\overline{T}_{\text{sing}})^- = \overline{\text{dom } T} \times \text{mul } \overline{T}.$$

Proof. By Proposition 4.5 the lefthand side is equal to $(T_{\text{sing}})^-$. Moreover,

$$(T_{\text{sing}})^- = \overline{\text{dom } T_{\text{sing}}} \times \text{mul } (T_{\text{sing}})^- = \overline{\text{dom } T} \times \text{mul } \overline{T},$$

due to Proposition 3.2 and Theorem 4.1. $\square$

Remark 4.7. The relation $\overline{T}_{\text{reg}}$ is closed; however, in general, the relation $\overline{T}_{\text{sing}}$ is not closed. For assume that $\overline{T}_{\text{sing}}$ is closed. Then by Corollary 4.6

$$\overline{T}_{\text{sing}} = \overline{\text{dom } T} \times \text{mul } \overline{T}.$$

This implies that $\text{dom } \overline{T}_{\text{sing}} = \overline{\text{dom } T}$, whereas by definition $\text{dom } \overline{T}_{\text{sing}} = \text{dom } \overline{T}$. Hence, if $\overline{T}_{\text{sing}}$ is closed, then $\text{dom } \overline{T}$ is closed.

5. A METRIC CHARACTERIZATION

Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$, and introduce the graph inner product on T by:

$$(\{f, f'\}, \{g, g'\})_T \stackrel{\text{def}}{=} (f, g) + (f', g'), \quad \{f, f'\}, \{g, g'\} \in T.$$

Denote the corresponding inner product space by $\mathfrak{G}(T)$. Define the mapping ι_T by

$$\iota_T \{f, f'\} \stackrel{\text{def}}{=} f, \quad \{f, f'\} \in T,$$

so that ι_T is a contraction on $\mathfrak{G}(T)$ with values in $\text{dom } T \subset \mathfrak{H}$. The isometric part of ι_T is its restriction to $\ker T \times \{0\}$. Note that the Hilbert space completion of $\mathfrak{G}(T)$ is $\mathfrak{G}(\overline{T})$, where $\mathfrak{G}(\overline{T})$ stands for the construction related to the closure $\overline{T}$ of T . Clearly $\mathfrak{G}(\overline{T}) = \mathfrak{G}(T)$ if and only if the relation T is closed, in which case the contraction ι_T is closed. In general, the contraction ι_T has a closure $\overline{\iota}_T$, which is a contraction defined on all of $\overline{T}$. The contraction $\iota_{\overline{T}}$ is also defined everywhere on $\overline{T}$. Clearly the restrictions of the contractions $\overline{\iota}_T$ and $\iota_{\overline{T}}$ coincide on T , which is a dense set in $\overline{T}$; hence they coincide:

$$\overline{\iota}_T = \iota_{\overline{T}}.$$

It follows from $\text{dom } T \subset \text{dom } \overline{T} \subset \overline{\text{dom } T}$ that $\overline{\iota}_T = \iota_{\overline{T}}$ is a contraction defined on all of $\mathfrak{G}(\overline{T})$ with values in $\overline{\text{dom } T}$. Since $\iota_{\overline{T}}$ maps $\mathfrak{G}(\overline{T})$ into $\overline{\text{dom } T}$, it follows by the first identity in (2.2) that

$$\overline{\text{ran}}(\iota_T)^* = \overline{\text{ran}}(\iota_{\overline{T}})^* = \mathfrak{G}(\overline{T}) \ominus \ker \iota_{\overline{T}}.$$

In particular, this shows that

$$(5.1) \quad (\iota_T)^* \upharpoonright \text{dom } T \text{ is dense in } \mathfrak{G}(\overline{T}) \ominus \ker \iota_{\overline{T}}.$$

The above construction is the analog for relations of a construction involving semibounded sesquilinear forms, cf. [9], [26]. The following lemma is obvious.

Lemma 5.1. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then*

$$(5.2) \quad \ker \iota_{\overline{T}} = \{0\} \times \text{mul } \overline{T}.$$

Denote by Q the orthogonal projection from the Hilbert space $\mathfrak{G}(\overline{T})$ onto its closed linear subspace $\ker \iota_{\overline{T}}$.

Lemma 5.2. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. The orthogonal projection Q in $\mathfrak{G}(\overline{T})$ and the orthogonal projection P in $\mathfrak{K}$ are related by*

$$Q\{\varphi, \varphi'\} = \{0, P\varphi'\}, \quad (I - Q)\{\varphi, \varphi'\} = \{\varphi, (I - P)\varphi'\}, \quad \{\varphi, \varphi'\} \in T.$$

Proof. In the sense of the inner product of the Hilbert space $\mathfrak{G}(\overline{T})$ each element $\{\varphi, \varphi'\} \in T$ has the orthogonal decomposition

$$(5.3) \quad \{\varphi, \varphi'\} = \{h, h'\} \hat{+} \{k, k'\},$$

where

$$(5.4) \quad \{h, h'\} = (I - Q)\{\varphi, \varphi'\}, \quad \{k, k'\} = Q\{\varphi, \varphi'\}.$$

The elements $\{h, h'\}$ and $\{k, k'\}$ are orthogonal in the sense of $\mathfrak{G}(\overline{T})$:

$$0 = (h, k) + (h', k').$$

It follows from Lemma 5.1 that $k = 0$, so that $h = \varphi$ and $(h', k') = 0$. Hence the decomposition (5.3) can also be written as

$$(5.5) \quad \{\varphi, \varphi'\} = \{\varphi, h' + k'\}, \quad k' \in \text{mul } \overline{T},$$

where the decomposition of $\varphi' = h' + k'$ is orthogonal in $\mathfrak{H}$. In other words

$$h' = (I - P)\varphi', \quad k' = P\varphi',$$

so that (5.4) reads as

$$(I - Q)\{\varphi, \varphi'\} = \{\varphi, (I - P)\varphi'\}, \quad Q\{\varphi, \varphi'\} = \{0, P\varphi'\},$$

which gives the statement of the lemma. $\square$

The construction involving the contraction ι_T now leads to a metric characterization of the elements in the ranges of T_{reg} and T_{sing} .

Lemma 5.3. *Let T be a linear relation from the Hilbert space $\mathfrak{H}$ to the Hilbert space $\mathfrak{K}$. Then for all $\{\varphi, \varphi'\} \in T$:*

$$(5.6) \quad \|P\varphi'\|^2 = \|\varphi'\|^2 + \inf_{h \in \text{dom } T} \left\{ \|\varphi + h\|^2 - \inf_{\{g, g'\} \in T} \{\|g'\|^2 + \|g + h\|^2\} \right\},$$

and

$$(5.7) \quad \|(I - P)\varphi'\|^2 = \sup_{h \in \text{dom } T} \inf_{\{g, g'\} \in T} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|g'\|^2 \}$$

Proof. According to Lemma 5.2 $Q\{\varphi, \varphi'\} = \{0, P\varphi'\}$, which implies that

$$(5.8) \quad \|P\varphi'\|^2 = \|Q\{\varphi, \varphi'\}\|_{\overline{T}}^2.$$

Recall that $\mathfrak{G}(\overline{T}) = \ker \iota_{\overline{T}} \oplus (\mathfrak{G}(\overline{T}) \ominus \ker \iota_{\overline{T}})$, which gives

$$\|Q\{\varphi, \varphi'\}\|_{\overline{T}}^2 = \inf \{ \|\{\varphi, \varphi'\} - \{\alpha, \alpha'\}\|_{\overline{T}}^2 : \{\alpha, \alpha'\} \in \mathfrak{G}(\overline{T}) \ominus \ker \iota_{\overline{T}} \}.$$

However, since $\iota_T^*(\text{dom } T)$ is dense in $\mathfrak{G}(\overline{T}) \ominus \ker \iota_{\overline{T}}$, it follows that

$$(5.9) \quad \begin{aligned} \|Q\{\varphi, \varphi'\}\|_{\overline{T}}^2 &= \inf_{h \in \text{dom } T} \{ \|\{\varphi, \varphi'\} + \iota_T^* h, \{\varphi, \varphi'\} + \iota_T^* h\|_{\overline{T}}^2 \\ &= \inf_{h \in \text{dom } T} \{ (\varphi, \varphi) + (\varphi', \varphi') + (\varphi, h) + (h, \varphi) + (\iota_T^* h, \iota_T^* h)_{\overline{T}} \} \\ &= \|\varphi'\|^2 + \inf_{h \in \text{dom } T} \{ \|\varphi + h\|^2 - \|h\|^2 + \|\iota_T^* h\|_{\overline{T}}^2 \}. \end{aligned}$$

Furthermore, since T is dense in $\mathfrak{G}(\overline{T})$, every element of the form $\iota_T^* h$, $h \in \text{dom } T$, can be approximated by elements in T , which leads to

$$(5.10) \quad \begin{aligned} 0 &= \inf_{\{g, g'\} \in T} \{ \|\{g, g'\} + \iota_T^* h\|_{\overline{T}}^2 \} \\ &= \|\iota_T^* h\|_{\overline{T}}^2 + \inf_{\{g, g'\} \in T} \{ (g, g) + (g', g') + (g, h) + (h, g) \} \\ &= -\|h\|^2 + \|\iota_T^* h\|_{\overline{T}}^2 + \inf_{\{g, g'\} \in T} \{ \|g'\|^2 + \|g + h\|^2 \}. \end{aligned}$$

Combining the identities (5.9), (5.10), and (5.8) gives (5.6). Clearly (5.7) follows from (5.6). $\square$

The above lemma leads to a metric characterization of closable operators.

Theorem 5.4. *Let T be a linear operator from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then T is closable if and only if for all $\varphi \in \text{dom } T$:*

$$(5.11) \quad \|T\varphi\|^2 = \sup_{h \in \text{dom } T} \inf_{g \in \text{dom } T} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Tg\|^2 \}.$$

Proof. Assume that (5.11) holds for all $\varphi \in \text{dom } T$. By Lemma 5.3 this means that

$$\|T\varphi\|^2 = \|(I - P)T\varphi\|^2, \quad \varphi \in \text{dom } T.$$

In other words, $PT\varphi = 0$ and hence $T\varphi \in \ker P$ for all $\varphi \in \text{dom } T$. Therefore $\text{ran } T \subset \ker P$, which leads to

$$\text{ran } \overline{T} \subset \ker P, \quad \text{in particular,} \quad \text{mul } \overline{T} \subset \ker P.$$

Since P is an orthogonal projection onto $\text{mul } \overline{T}$, this implies that $\text{mul } \overline{T} = \{0\}$, i.e., $\overline{T}$ is an operator. Hence, the operator T is closable.

Conversely, assume that the operator T is closable. Then according to Theorem 4.1 $T = T_{\text{reg}}$ and $P = 0$. Hence if $\{\varphi, \varphi'\} \in T$, then $T\varphi = \varphi' = (I - P)\varphi'$ and the result follows from (5.7). $\square$

In general the supremum and the infimum in (5.11) are not attained. However, when the operator T is densely defined and closed one can say more.

Lemma 5.5. *Let T be a densely defined closed linear operator from $\mathfrak{H}$ to $\mathfrak{K}$. Then*

$$(5.12) \quad \min_{g \in \text{dom } T} (\|g + h\|^2 + \|Tg\|^2) = \|h\|^2 - \|(I + T^*T)^{-\frac{1}{2}}h\|^2, \quad h \in \mathfrak{H},$$

*and the minimum is attained for $g = -(I + T^*T)^{-1}h \in \text{dom } T^*T$.*

Proof. Since T is densely defined and closed, T^*T is a selfadjoint operator, which is nonnegative. Observe that for all $g \in \text{dom } T$ there is the identity

$$\|g\|^2 + \|Tg\|^2 = \|(I + T^*T)^{\frac{1}{2}}g\|^2.$$

Hence, for all $g \in \text{dom } T$ and $h \in \mathfrak{H}$ one has

$$\begin{aligned} & \|g + h\|^2 + \|Tg\|^2 \\ &= \|h\|^2 - \|(I + T^*T)^{-\frac{1}{2}}h\|^2 + \left\| (I + T^*T)^{\frac{1}{2}}g + (I + T^*T)^{-\frac{1}{2}}h \right\|^2 \\ &\geq \|h\|^2 - \|(I + T^*T)^{-\frac{1}{2}}h\|^2. \end{aligned}$$

Due to $\text{ran } (I + T^*T)^{\frac{1}{2}} = \mathfrak{H}$ this implies that (5.12) holds and the minimum is attained for $g = -(I + T^*T)^{-1}h \in \text{dom } T^*T$. $\square$

A combination of Theorem 5.4 and Lemma 5.5 leads to the following characterization.

Proposition 5.6. *Let T be a densely defined closed linear operator from $\mathfrak{H}$ to $\mathfrak{K}$. Then*

$$(5.13) \quad \|T\varphi\|^2 = \sup_{h \in \text{dom } T} \min_{g \in \text{dom } T} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Tg\|^2 \},$$

*and the supremum is a maximum if and only if $\varphi \in \text{dom } T^*T$.*

Proof. Since the operator T is closed the identity (5.11) holds for all $\varphi \in \text{dom } T$. Clearly, it follows from (5.11) and (5.12) that

$$(5.14) \quad \|T\varphi\|^2 = \sup_{h \in \text{dom } T} \left(\|h\|^2 - \|(I + T^*T)^{-\frac{1}{2}}h\|^2 - \|\varphi + h\|^2 \right), \quad \varphi \in \text{dom } T.$$

Observe that for all $\varphi \in \text{dom } T$ and $h \in \mathfrak{H}$ one has

$$\begin{aligned} & \|h\|^2 - \|(I + T^*T)^{-\frac{1}{2}}h\|^2 - \|\varphi + h\|^2 \\ &= -\|(I + T^*T)^{-\frac{1}{2}}h + (I + T^*T)^{\frac{1}{2}}\varphi\|^2 + \|T\varphi\|^2, \end{aligned}$$

which implies that the supremum in (5.14) and hence in (5.13) is a maximum if and only if $\varphi \in \text{dom } T^*T$. $\square$

In particular, if T is a bounded linear operator from $\mathfrak{H}$ to $\mathfrak{K}$ then the supremum in (5.13) can be replaced by a maximum. The original observation about the minimum and maximum in (5.11) for bounded linear operators goes back to L. László (personal communication, 2005).

Corollary 5.7. *Let T be a closed linear operator from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ and let S be a linear operator from the Hilbert space $\mathfrak{H}$ to the Hilbert space $\mathfrak{K}$ with $\text{dom } S = \text{dom } T$ and*

$$(5.15) \quad \|T\varphi\| \leq \|S\varphi\|, \quad \varphi \in \mathfrak{D} \stackrel{\text{def}}{=} \text{dom } S = \text{dom } T.$$

Then for all $\varphi \in \mathfrak{D}$:

$$(5.16) \quad \|T\varphi\|^2 \leq \sup_{h \in \mathfrak{D}} \inf_{g \in \mathfrak{D}} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Sg\|^2 \} \leq \|S\varphi\|^2.$$

Proof. It follows from Theorem 5.4 and the assumption (5.15) that

$$\begin{aligned} \|T\varphi\|^2 &= \sup_{h \in \mathfrak{D}} \inf_{g \in \mathfrak{D}} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Tg\|^2 \} \\ &\leq \sup_{h \in \mathfrak{D}} \inf_{g \in \mathfrak{D}} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Sg\|^2 \}, \quad \varphi \in \mathfrak{D}. \end{aligned}$$

The choice $g = \varphi$ leads to the inequality

$$\sup_{h \in \mathfrak{D}} \inf_{g \in \mathfrak{D}} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Sg\|^2 \} \leq \|S\varphi\|^2,$$

which results in (5.16). $\square$

The result in Theorem 5.4 can be interpreted in terms of parallel sums and differences (see [9]), cf. [1], [2], [18], [29]. Furthermore, by replacing T with its inverse T^{-1} one obtains a similar characterization for the implication $\ker T = \{0\} \Rightarrow \ker \overline{T} = \{0\}$. Observe also that if the operator T in Theorem 5.4 is not closable, then for some $\varphi \in \text{dom } T$:

$$\|T\varphi\|^2 > \sup_{h \in \text{dom } T} \inf_{g \in \text{dom } T} \{ \|g + h\|^2 - \|\varphi + h\|^2 + \|Tg\|^2 \}.$$

6. THE STONE DECOMPOSITION

Since a closed linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ is by definition a closed linear subspace of the Cartesian product $\mathfrak{H} \times \mathfrak{K}$, there is a one-to-one correspondence between the orthogonal projections in $\mathfrak{H} \times \mathfrak{K}$ and the closed linear relations from $\mathfrak{H}$ to $\mathfrak{K}$. This section gives a short review of the consequences of this correspondence, cf. [19], [28].

Let $\mathfrak{H}$ and $\mathfrak{K}$ be Hilbert spaces and let R be an orthogonal projection on the Cartesian product $\mathfrak{H} \times \mathfrak{K}$. Decompose R according to the Cartesian product

$$(6.1) \quad R = \begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix},$$

so that $R_{11} \geq 0$, $R_{22} \geq 0$, and $R_{21} = R_{12}^*$.

Lemma 6.1 ([19]). *The entries in the block decomposition (6.1) satisfy:*

- (i) $\ker R_{11} \subset \ker R_{21}$;
- (ii) $\ker R_{22} \subset \ker R_{12}$;
- (iii) $\ker (I - R_{11}) \subset \ker R_{21}$;
- (iv) $\ker (I - R_{22}) \subset \ker R_{12}$.

Let T be a closed relation from $\mathfrak{H}$ to $\mathfrak{K}$ and let R be the corresponding orthogonal projection onto T . The corresponding matrix $R = (R_{ij})$ is called the *characteristic matrix* of T . It follows by definition that

$$(6.2) \quad T = \{ \{ R_{11}h + R_{12}h', R_{21}h + R_{22}h' \} : \{ h, h' \} \in \mathfrak{H} \times \mathfrak{K} \}.$$

The representation (6.2) is called the *Stone decomposition* of T . It is clear that

$$(6.3) \quad \begin{pmatrix} R_{22} & R_{21} \\ R_{12} & R_{11} \end{pmatrix}$$

is the characteristic matrix of the inverse relation T^{-1} (keeping the order of the parametrizing elements in mind).

Lemma 6.2. *Let T be a closed linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$, and let R be its characteristic matrix. Then*

$$(6.4) \quad \text{dom } T = \text{ran} \begin{pmatrix} R_{11} & R_{12} \end{pmatrix}, \quad \text{ran } T = \text{ran} \begin{pmatrix} R_{21} & R_{22} \end{pmatrix},$$

and

$$(6.5) \quad \ker T = \ker (I - R_{11}), \quad \text{mul } T = \ker (I - R_{22}).$$

Proof. The identities in (6.4) are clear. Observe also that $\text{ran } T = \text{dom } T^{-1}$, so that, in fact, the second identity in (6.4) follows from the first identity and (6.3).

To prove the first identity in (6.5), let $h \in \ker T$, then $\{h, 0\} \in T$, so that in particular $\{h, 0\} = R\{h, 0\}$, or equivalently,

$$\begin{pmatrix} h \\ 0 \end{pmatrix} = \begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \begin{pmatrix} h \\ 0 \end{pmatrix} = \begin{pmatrix} R_{11}h \\ R_{21}h \end{pmatrix},$$

which implies that $R_{11}h = h$ and $R_{21}h = 0$, and, by Lemma 6.1, this is equivalent to $h \in \ker (I - R_{11})$.

The statement concerning $\text{mul } T = \ker T^{-1}$ follows from the first identity in (6.5) and (6.3). $\square$

Corollary 6.3. *Let T be a closed linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Its regular part T_{reg} is given by*

$$(6.6) \quad T_{\text{reg}} = \{ \{R_{11}h + R_{12}k, R_{21}h + R_{22}k\} : h \in \mathfrak{H}, k \in (\text{mul } T)^\perp \}.$$

Proof. Consider the representation (6.2) of the closed linear relation T . Decompose the variable $h' \in \mathfrak{H}$ by $h' = k + \varphi$ with $k \in (\text{mul } T)^\perp$ and $\varphi \in \text{mul } T$. Then it follows from Lemma 6.1 and Lemma 6.2 that

$$R_{11}h + R_{12}h' = R_{11}h + R_{12}k, \quad R_{21}h + R_{22}h' = R_{21}h + R_{22}k + \varphi,$$

and it is also clear that $R_{21}h + R_{22}k \in (\text{mul } T)^\perp$. This completes the proof. $\square$

The definition of the adjoint $T^* = JT^\perp$ (where the product in the righthand side is carried out in the indicated order) gives the corresponding characteristic matrix

$$(6.7) \quad \begin{pmatrix} I - R_{22} & R_{21} \\ R_{12} & I - R_{11} \end{pmatrix},$$

which leads to the following parametrization of T^* :

$$(6.8) \quad T^* = \{ \{(I - R_{22})h + R_{21}h', R_{12}h + (I - R_{11})h'\} : \{h, h'\} \in \mathfrak{H} \times \mathfrak{K} \}.$$

By Lemma 6.2 it follows that

$$(6.9) \quad \text{dom } T^* = \text{ran} \begin{pmatrix} (I - R_{22}) & R_{21} \end{pmatrix}, \quad \text{ran } T^* = \text{ran} \begin{pmatrix} R_{12} & (I - R_{11}) \end{pmatrix},$$

and

$$(6.10) \quad \ker T^* = \ker R_{22}, \quad \text{mul } T^* = \ker R_{11}.$$

The regular part of T^* is given by

$$(T^*)_{\text{reg}} = \{ \{(I - R_{22})h + R_{21}k, R_{12}h + (I - R_{11})k\} : h \in \mathfrak{H}, k \in (\text{mul } T^*)^\perp \}.$$

The following result is now straightforward, see [28] for the case of operators and [16] for the case of relations.

Lemma 6.4. *Let T be a closed linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then the products T^*T and TT^* are nonnegative selfadjoint relations in $\mathfrak{H}$ and $\mathfrak{K}$, respectively, and*

$$(T^*T + I)^{-1} \quad \text{and} \quad (TT^* + I)^{-1}$$

are (the graphs of) bounded linear operators defined on all of $\mathfrak{H}$ and $\mathfrak{K}$, respectively. The characteristic matrix R of T is given by

$$R = \begin{pmatrix} (T^*T + I)^{-1} & (T^*)_{\text{reg}}(TT^* + I)^{-1} \\ T_{\text{reg}}(T^*T + I)^{-1} & I - (TT^* + I)^{-1} \end{pmatrix}.$$

Proof. Let $h \in \mathfrak{H}$, then there is a unique decomposition

$$\{h, 0\} = \{\varphi, \varphi'\} + \{\psi, \psi'\}, \quad \{\varphi, \varphi'\} \in T, \quad \{\psi, \psi'\} \in T^\perp = JT^*,$$

since $\mathfrak{H} \times \mathfrak{K} = T \hat{\oplus} T^\perp$. Hence

$$h = \varphi + \psi, \quad \varphi' + \psi' = 0,$$

which leads to $\{\psi, \psi'\} = \{\psi, -\varphi'\} \in JT^*$ and $\{\varphi', \psi\} \in T^*$. Therefore, $\{\varphi, \psi\} \in T^*T$ and

$$\{\varphi, h\} = \{\varphi, \varphi + \psi\} \in T^*T + I,$$

so that $h \in \text{ran}(T^*T + I)$. Thus $\text{ran}(T^*T + I) = \mathfrak{H}$ and $\varphi = (T^*T + I)^{-1}h$. Note that $\{\psi, \psi'\} \in T^\perp = JT^*$ or, equivalently, $\{\psi', -\psi\} \in T^*$, so that $\varphi' = -\psi' \in \text{dom } T^* \subset (\text{mul } T)^\perp$ and it follows that $\varphi' = T_{\text{reg}}\varphi$. Therefore

$$\varphi = (T^*T + I)^{-1}h, \quad \varphi' = T_{\text{reg}}(T^*T + I)^{-1}h.$$

Hence the first column of R is completely determined. By formally replacing T by T^* also the first column of the characteristic matrix of T^* is determined. Now the second column of R is obtained via (6.7). $\square$

The statements concerning the products T^*T and TT^* are known (cf. [7]), but the proof is repeated here to make the identification of the first column of R understandable.

7. A CHARACTERIZATION OF THE REGULAR AND SINGULAR PARTS VIA THE STONE DECOMPOSITION

Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ and let $\overline{T}$ be its closure. Decompose the Cartesian product $\mathfrak{H} \times \mathfrak{K}$ as follows

$$(7.1) \quad \mathfrak{H} \times \mathfrak{K} = \overline{T} \hat{\oplus} T^\perp,$$

where $\hat{\oplus}$ denotes the orthogonal component-wise sum. Let $\tilde{R}$ denote the orthogonal projection from $\mathfrak{H} \times \mathfrak{K}$ onto $\overline{T}$ and decompose $\tilde{R}$ according to the Cartesian product:

$$\tilde{R} = \begin{pmatrix} \tilde{R}_{11} & \tilde{R}_{12} \\ \tilde{R}_{21} & \tilde{R}_{22} \end{pmatrix}.$$

Observe that

$$(7.2) \quad \text{mul } \overline{T} = \ker(I - \tilde{R}_{22}) \subset \ker \tilde{R}_{12},$$

by combining Lemma 6.2 and Lemma 6.1, and that

$$(7.3) \quad \text{mul } T^* = \ker \tilde{R}_{11} \subset \ker \tilde{R}_{21},$$

by combining (6.10) and Lemma 6.1. In particular, this implies that

$$(7.4) \quad \tilde{R}_{11} \upharpoonright_{\text{mul } T^*} = 0 \upharpoonright_{\text{mul } T^*}, \quad \tilde{R}_{21} \upharpoonright_{\text{mul } T^*} = 0 \upharpoonright_{\text{mul } T^*},$$

and

$$(7.5) \quad \tilde{R}_{12} \upharpoonright_{\text{mul } \overline{T}} = 0 \upharpoonright_{\text{mul } \overline{T}}, \quad \tilde{R}_{22} \upharpoonright_{\text{mul } \overline{T}} = I \upharpoonright_{\text{mul } \overline{T}}.$$

Proposition 7.1. *The relation T is regular if and only if*

$$(7.6) \quad \ker (I - \tilde{R}_{22}) = \{0\}.$$

Proof. By definition the relation T is regular if and only if $\overline{T}$ is an operator, or, equivalently, $\text{mul } \overline{T} = \{0\}$. Hence the identity (7.6) follows from the first identity in (7.2). $\square$

Proposition 7.2. *The relation T is singular if and only if*

$$(7.7) \quad \tilde{R}_{12} = 0 \quad \text{or equivalently} \quad \tilde{R}_{21} = 0,$$

in which case

$$(7.8) \quad \tilde{R} = \begin{pmatrix} I \upharpoonright_{\overline{\text{dom}} T} \oplus 0 \upharpoonright_{(\text{dom } T)^\perp} & 0 \\ 0 & I \upharpoonright_{\overline{\text{ran}} T} \oplus 0 \upharpoonright_{(\text{ran } T)^\perp} \end{pmatrix}.$$

Proof. Let T be a singular relation. Then $\text{ran } T \subset \text{mul } \overline{T}$ and it follows from (7.2) that for all $\{\varphi, \varphi'\} \in T \subset \overline{T}$

$$\begin{pmatrix} \varphi \\ \varphi' \end{pmatrix} = \begin{pmatrix} \tilde{R}_{11} & \tilde{R}_{12} \\ \tilde{R}_{21} & \tilde{R}_{22} \end{pmatrix} \begin{pmatrix} \varphi \\ \varphi' \end{pmatrix} = \begin{pmatrix} \tilde{R}_{11}\varphi \\ \tilde{R}_{21}\varphi + \varphi' \end{pmatrix},$$

as $\varphi' \in \text{mul } \overline{T}$. Hence for all $\varphi \in \text{dom } T$:

$$(7.9) \quad \tilde{R}_{11}\varphi = \varphi \quad \text{and} \quad \tilde{R}_{21}\varphi = 0.$$

Since the entries in the block decomposition of $\tilde{R}$ are bounded operators, one has $\tilde{R}_{21} \upharpoonright_{\overline{\text{dom}} T} = 0 \upharpoonright_{\overline{\text{dom}} T}$, which together with the second identity in (7.4) shows that $\tilde{R}_{21} = 0$. This gives (7.7).

Conversely, assume that (7.7) holds. Then necessarily $\tilde{R}_{22}$ is an orthogonal projection in $\mathfrak{K}$ and now Lemma 6.2 gives

$$(7.10) \quad \text{ran } T \subset \text{ran } \overline{T} = \text{ran } \tilde{R}_{22} = \ker (I - \tilde{R}_{22}) = \text{mul } \overline{T},$$

so that T is singular.

It remains to discuss the matrix representation (7.8). The representation of $\tilde{R}_{11}$ is obtained from the first identities in (7.4) and (7.9). The representation of $\tilde{R}_{22}$ follows immediately from (7.10), since $\tilde{R}_{22}$ is an orthogonal projection in $\mathfrak{K}$. $\square$

The result in Proposition 7.2 shows that there is a one-to-one correspondence between the closed singular relations from $\mathfrak{H}$ to $\mathfrak{K}$ and the orthogonal projections R in $\mathfrak{H} \times \mathfrak{K}$ which are of the form $R = R_{11} \oplus R_{22}$, where R_{11} is an orthogonal projection in $\mathfrak{H}$ and R_{22} is an orthogonal projection in $\mathfrak{K}$. Maximally singular relations can be described as follows.

Corollary 7.3. *The relation T is maximally singular if and only if*

$$(7.11) \quad \tilde{R}_{22} = I,$$

in which case (7.7) holds and

$$(7.12) \quad \tilde{R} = \begin{pmatrix} I \upharpoonright_{\overline{\text{dom}} T} \oplus 0 \upharpoonright_{(\text{dom } T)^\perp} & 0 \\ 0 & I \end{pmatrix}.$$

Proof. By Lemma 6.2 (see (7.2)) $\text{mul } \overline{T} = \ker (I - \tilde{R}_{22})$ and thus T is maximally singular, i.e. $\text{mul } \overline{T} = \mathfrak{K}$, if and only if $\tilde{R}_{22} = I$. Observe that (7.11) implies (7.7) in view of (7.2).

The representation (7.12) is immediate from (7.8). $\square$

8. EXAMPLES OF CANONICAL DECOMPOSITIONS

This section contains some illustrative examples concerning the canonical decomposition of (not necessarily densely defined) linear operators and relations.

Example 8.1. [27] Let S be a closable operator in a Hilbert space $\mathfrak{H}$ and let $Z = \text{span}\{e, f\}$ be a one-dimensional subspace in the Cartesian product $\mathfrak{H} \times \mathfrak{H}$. Note that $\{e, f\} \in S$ if and only if $e \in \text{dom } S$ and $Se = f$. Hence the assumption $S \cap Z = \{0, 0\}$ is equivalent to the assumption that either $e \in \text{dom } S$ and $Se \neq f$ or $e \notin \text{dom } S$. Under this assumption the component-wise sum

$$(8.1) \quad A = S \hat{+} Z,$$

defines a one-dimensional extension (of the graph) of S . It follows from (8.1) that $A^{**} = S^{**} \hat{+} Z$, since Z is one-dimensional. The one-dimensional extension A is not regular if and only if there is a nontrivial $k \in \mathfrak{H}$ for which $\{0, k\} \in \overline{A} = A^{**}$, i.e.,

$$\{0, k\} = \{h, \overline{S}h\} \hat{+} \lambda\{e, f\},$$

for some $h \in \text{dom } \overline{S} = \text{dom } S^{**}$ and $\lambda \in \mathbb{C}$. Equivalently, the one-dimensional extension A is not regular if and only if

$$(8.2) \quad e \in \text{dom } \overline{S}, \quad \overline{S}e \neq f,$$

in which case

$$\text{mul } \overline{A} = \text{span}\{\overline{S}e - f\}.$$

According to the assumption that the sum (8.1) is direct there are two cases for the relation A not to be regular. In the first case $e \in \text{dom } S$ and $Se \neq f$ and (8.2) is satisfied; this actually means that $\text{mul } A$ is not trivial and A itself is not an operator. In the second case $e \notin \text{dom } S$ and (8.2) means $e \in \text{dom } \overline{S} \setminus \text{dom } S$ and $\overline{S}e \neq f$; in this case A is an operator and $\text{mul } \overline{A}$ is one-dimensional.

In other words, under the assumption that the sum (8.1) is direct, the relation A is regular if and only if $e \in \text{dom } \overline{S} \setminus \text{dom } S$ and $\overline{S}e = f$, or $e \notin \text{dom } \overline{S}$.

Example 8.2. Let A be an operator in a Hilbert space $\mathfrak{H}$ which is not closable with $\text{mul } \overline{A} = \text{span}\{\varphi\}$, $\|\varphi\| = 1$, cf. Example 8.1. Denote the orthogonal projection onto $\text{mul } \overline{A}$ by P . Then

$$Ah = (I - P)Ah + PAh, \quad h \in \text{dom } A,$$

and

$$PAh = (PAh, \varphi)\varphi = (Ah, \varphi)\varphi, \quad h \in \text{dom } A.$$

Hence, by definition,

$$A_{\text{reg}}h = Ah - (Ah, \varphi)\varphi, \quad A_{\text{sing}}h = (Ah, \varphi)\varphi, \quad h \in \text{dom } A,$$

and by Theorem 4.1 A_{reg} is regular and A_{sing} is singular. According to Proposition 3.2 one has

$$(A_{\text{sing}})^- = \overline{\text{dom } A} \times \text{span}\{\varphi\}.$$

Example 8.3. Let A be an operator which is not closable with $\text{mul } \overline{A} = \text{span}\{\varphi\}$, $\|\varphi\| = 1$, cf. Example 8.1, and let B be a bounded everywhere defined operator. Define the operator T as an operator sum by

$$T = A + B, \quad \text{dom } T = \text{dom } A.$$

Then $T^* = A^* + B^*$ and

$$\overline{T} = \overline{A} + B, \quad \text{dom } \overline{T} = \text{dom } \overline{A}.$$

Moreover, $\text{mul } \overline{T} = \text{mul } \overline{A} = \text{span}\{\varphi\}$, which implies that the operator $T = A + B$ is not closable. It follows from (4.1) and (4.2) that

$$T_{\text{reg}}h = [A_{\text{reg}} + (I - P)B]h, \quad T_{\text{sing}}h = A_{\text{sing}}h + (Bh, \varphi)\varphi, \quad h \in \text{dom } A.$$

Again from Proposition 3.2 one obtains

$$(T_{\text{sing}})^- = \overline{\text{dom } A} \times \text{span } \{\varphi\}.$$

In particular, if the operator A is singular, then its perturbation by the bounded operator $B \neq 0$ is singular if and only if $\text{ran } B = \text{span } \{\varphi\}$.

Example 8.4. Let A be a bounded operator in a Hilbert space $\mathfrak{H}$ with domain $\text{dom } A$, let $\mathfrak{R}$ be a not necessarily closed subspace of $\mathfrak{H}$, and define the linear relation T by

$$T = \{ \{f, Af + \varphi\} : f \in \text{dom } A, \varphi \in \mathfrak{R} \}.$$

It is not difficult to see that

$$\overline{T} = \{ \{f, \overline{A}f + \varphi\} : f \in \text{dom } \overline{A}, \varphi \in \mathfrak{R}^- \},$$

since $\overline{A}$ is a bounded operator, cf. (2.4) and (2.5). Now this identity implies that $\text{mul } \overline{T} = \mathfrak{R}^-$. Hence

$$T_{\text{reg}} = \{ \{f, (I - P)Af\} : f \in \text{dom } A \},$$

and

$$T_{\text{sing}} = \{ \{f, PAf + \varphi\} : f \in \text{dom } A, \varphi \in \mathfrak{R} \}.$$

It follows from Proposition 3.2 that

$$(T_{\text{sing}})^- = \overline{\text{dom } A} \times \mathfrak{R}^-.$$

Note that T is maximally singular if and only if $\mathfrak{R}$ is dense in $\mathfrak{H}$. If in this example A is the null operator on $\mathfrak{D} = \text{dom } A$, then T reduces to the Cartesian product

$$T = \mathfrak{D} \times \mathfrak{R},$$

which is a singular relation.

9. AN APPROACH VIA ADJOINT RELATIONS

There is another approach to the decomposition results in Section 4. It is based on a description of the domains of the adjoints of T_{reg} and T_{sing} . First a general characterization result will be described.

Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. The following lemma gives a description of $\text{dom } T^*$. Clearly, by inverting the linear relation, a similar result can be obtained for the description of $\text{ran } T^*$. This latter description goes back to [24] in the case of bounded linear operators (see also [5]) and to [22] for densely defined operators; the present version can be found in [6], the proof is presented here for completeness. The orthogonal decomposition of a closed linear relation K into an operator part and a multivalued part is already assumed here (see Proposition 4.4):

$$(9.1) \quad K = K_s \hat{\oplus} K_\infty,$$

where $K_\infty \stackrel{\text{def}}{=} \{0\} \times \text{mul } K$ and $K_s \stackrel{\text{def}}{=} K \hat{\ominus} K_\infty (= K_{\text{reg}})$.

Lemma 9.1. ([6]) *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$. Then $g \in \text{dom } T^*$ if and only if there exists a nonnegative number C_g , such that*

$$(9.2) \quad |(f', g)_\mathfrak{K}| \leq C_g \|f\|_\mathfrak{H} \quad \text{for all } \{f, f'\} \in T.$$

In this case the smallest C_g satisfying (9.2) is $C_g = \|g'\|_\mathfrak{H}$ with $\{g, g'\} \in T^$ and $g' \in \text{dom } T$, i.e., $C_g = \|(T^*)_s g\|_\mathfrak{H}$.*

Proof. First assume that $g \in \text{dom } T^*$. Then $\{g, g'\} \in T^*$ for some $g' \in \mathfrak{H}$ and by the definition of the adjoint T^* one obtains for every $\{f, f'\} \in T$:

$$|(f', g)_\mathfrak{K}| = |(f, g')_\mathfrak{H}| \leq \|f\|_\mathfrak{H} \|g'\|_\mathfrak{H},$$

so that one can take $C_g = \|g'\|_\mathfrak{H}$ in (9.2).

Conversely, assume that $g \in \mathfrak{K}$ satisfies the estimate (9.2). Define the linear relation L_g in $\mathfrak{H} \oplus \mathbb{C}$ by

$$L_g := \{ \{f, (f', g)_{\mathfrak{K}}\} : \{f, f'\} \in T \}.$$

Then it follows from (9.2) that L_g is single-valued, since $\|f\|_{\mathfrak{H}} = 0$ implies $(f', g)_{\mathfrak{K}} = 0$. Hence, L_g is (the graph of) a single-valued bounded linear functional defined on $\text{dom } T$. Therefore, it has a continuation $\bar{L}_g$ from $\overline{\text{dom } T}$ into $\mathbb{C}$ with the same norm ($\|\bar{L}_g\| \leq C_g$). By the Riesz Representation Theorem there exists $g' \in \overline{\text{dom } T}$ with $\|g'\|_{\mathfrak{H}} = \|\bar{L}_g\|$, such that

$$\bar{L}_g f = (f, g')_{\mathfrak{H}} \quad \text{for all } f \in \overline{\text{dom } T}.$$

Therefore $(f', g)_{\mathfrak{K}} = (f, g')_{\mathfrak{H}}$ holds for every $\{f, f'\} \in T$, so that $\{g, g'\} \in T^*$. In particular, $g \in \text{dom } T^*$.

The last statement is clear from the given arguments and the definition of the orthogonal operator part $(T^*)_s$ of T^* , cf. (9.1). $\square$

Let T be a linear relation from a Hilbert space $\mathfrak{H}$ to a Hilbert space $\mathfrak{K}$ and let the regular and singular parts T_{reg} and T_{sing} of T be as defined in (4.1) and (4.2), respectively. In the present approach $\text{dom } (T_{\text{reg}})^*$, $\ker (T_{\text{reg}})^*$, $\text{dom } (T_{\text{sing}})^*$, and $\ker (T_{\text{sing}})^*$ are determined by means of Lemma 9.1.

Proposition 9.2. *Let T be a linear relation from a Hilbert space $\mathfrak{H}$ into a Hilbert space $\mathfrak{K}$. Then*

$$(9.3) \quad \text{dom } (T_{\text{reg}})^* = \text{dom } T^* \oplus \text{mul } \bar{T}, \quad \ker (T_{\text{reg}})^* = \ker T^* \oplus \text{mul } \bar{T},$$

and

$$(9.4) \quad \text{dom } (T_{\text{sing}})^* = \ker (T_{\text{sing}})^* = \overline{\text{dom } T^*}.$$

Proof. It follows from the definition of T_{reg} in (4.1) and from Lemma 9.1 that

$$(9.5) \quad g \in \text{dom } (T_{\text{reg}})^* \Leftrightarrow |(f', (I - P)g)| \leq C_g \|f\|, \quad \{f, f'\} \in T.$$

Clearly, the following equivalence is a consequence of (9.5):

$$g \in \text{dom } (T_{\text{reg}})^* \cap (\text{mul } \bar{T})^\perp \Leftrightarrow g \in \text{dom } T^* \cap (\text{mul } \bar{T})^\perp.$$

Therefore

$$\text{dom } (T_{\text{reg}})^* \cap \overline{\text{dom } T^*} = \overline{\text{dom } T^*} \cap \text{dom } T^*,$$

or, equivalently,

$$(9.6) \quad \text{dom } (T_{\text{reg}})^* \cap \overline{\text{dom } T^*} = \text{dom } T^*.$$

Furthermore it follows from (9.5) that

$$(9.7) \quad \text{mul } \bar{T} \subset \text{dom } (T_{\text{reg}})^*.$$

Since $\text{mul } \bar{T} \oplus \overline{\text{dom } T^*} = \mathfrak{K}$, the first identity in (9.3) is obtained from (9.6) and (9.7). In addition, by the second statement in Lemma 9.1 $g \in \ker T^* (= \ker (T^*)_s)$ if and only if the smallest constant in (9.2) is $C_g = 0$. Therefore, it follows from (9.5) that

$$(9.8) \quad \ker (T_{\text{reg}})^* \cap \overline{\text{dom } T^*} = \ker T^* \text{ and } \text{mul } \bar{T} \subset \ker (T_{\text{reg}})^*,$$

which proves the second identity in (9.3).

Likewise, it follows from the definition of T_{sing} in (4.2) and from Lemma 9.1 that

$$(9.9) \quad g \in \text{dom } (T_{\text{sing}})^* \Leftrightarrow |(f', Pg)| \leq C_g \|f\|, \quad \{f, f'\} \in T.$$

Clearly, the following equivalence is a consequence of (9.9):

$$g \in \text{dom } (T_{\text{sing}})^* \cap \text{mul } \bar{T} \Leftrightarrow g \in \text{dom } T^* \cap \text{mul } \bar{T}.$$

Observe that $\text{dom } T^* \cap \text{mul } \bar{T} = \{0\}$, which shows that

$$(9.10) \quad \text{dom } (T_{\text{sing}})^* \cap \text{mul } \bar{T} = \{0\}.$$

Furthermore it follows from (9.9) that

$$(9.11) \quad \overline{\text{dom}} T^* = (\text{mul } \overline{T})^\perp \subset \ker (T_{\text{sing}})^* \subset \text{dom} (T_{\text{sing}})^*$$

Since $\text{mul } \overline{T} \oplus \overline{\text{dom}} T^* = \mathfrak{K}$, the equalities in (9.4) follow from (9.10) and (9.11). $\square$

The results in Proposition 9.2 immediately give the main result in Section 4, see Theorem 4.1. For instance, according to (9.3)

$$\overline{\text{dom}} (T_{\text{reg}})^* = \overline{\text{dom}} T^* \oplus \text{mul } \overline{T} = \mathfrak{K},$$

and thus T_{reg} is regular by Proposition 3.1. Also the first equality in (9.4) shows that T_{sing} is singular by part (ii) of Proposition 3.2.

REFERENCES

- [1] T. Ando, Lebesgue-type decomposition of positive operators, *Acta Sci. Math. (Szeged)*, 38 (1976), 253–260.
- [2] T. Ando and W. Szymański, Order structure and Lebesgue decomposition of positive definite operator function, *Indiana University Math. J.*, 35 (1986), 157–173.
- [3] R. Arens, Operational calculus of linear relations, *Pacific J. Math.*, 9 (1961), 9–23.
- [4] E.A. Coddington, Extension theory of formally normal and symmetric subspaces, *Mem. Amer. Math. Soc.*, 134, 1973.
- [5] P. Fillmore and J. Williams, On operator ranges, *Adv. in Math.*, 7 (1971), 254–281.
- [6] S. Hassi, On the Friedrichs and the Krein-von Neumann extension of nonnegative relations, *Acta Wasaensia*, 122 (2004), 37–54.
- [7] S. Hassi, A. Sandovici, H.S.V. de Snoo, and H. Winkler, Form sums of nonnegative selfadjoint operators, *Acta Math. Hungar.*, 111 (1-2) (2006), 81–105.
- [8] S. Hassi, Z. Sebestyén, and H.S.V. de Snoo, On the nonnegativity of operator products, *Acta Math. Hungarica*, 109 (4) (2005), 1–15.
- [9] S. Hassi, Z. Sebestyén, and H.S.V. de Snoo, A canonical decomposition for a pair of nonnegative forms, in preparation.
- [10] S. Hassi, H.S.V. de Snoo, and F.H. Szafraniec, Normal extensions of symmetric relations, in preparation.
- [11] P.E.T. Jorgensen, Unbounded operators; perturbations and commutativity problems, *J. Functional Analysis*, 39 (1980), 281–307.
- [12] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1980.
- [13] H. Kosaki, Remarks on Lebesgue-type decomposition of positive operators, *J. Operator Theory*, 11 (1984), 137–143.
- [14] V.D. Koshmanenko, *Singular quadratic forms in perturbation theory*, Kluwer Academic Publishers, Dordrecht/ Boston/London, 1999.
- [15] V.D. Koshmanenko and S. Ôta, On characteristic properties of singular operators, *Ukr. Math. Zh.*, 48(1996), 1484–1493.
- [16] Y. Mezroui, Projection orthogonale sur le graphe d’une relation linéaire et fermé, *Trans. Amer. Math. Soc.*, 352(2000), 2789–2800.
- [17] J. von Neumann, *Functional operators*, Volume 2: The geometry of orthogonal spaces. *Ann. Math. Studies* No. 22, Princeton University Press, Princeton, N. J., 1950.
- [18] K. Nishio, Characterization of Lebesgue-type decomposition of positive operators, *Acta Sci. Math. (Szeged)*, 42(1980), 143–152.
- [19] A.E. Nussbaum, Reduction theory for unbounded closed operators in Hilbert space, *Duke Math. J.*, 31(1964), 33–44.
- [20] S. Ôta, Decomposition of unbounded derivations in operator algebras, *Tôhoku Math. J.*, 33(1981), 215–225.
- [21] S. Ôta, On a singular part of an unbounded operator, *Zeitschrift für Analysis und ihre Anwendungen*, 7(1987), 15–18.
- [22] Z. Sebestyén, On ranges of adjoint operators in Hilbert space, *Acta Sci. Math. (Szeged)*, 46 (1983), 295–298.
- [23] Z. Sebestyén, Operator extension, dilation, lifting, *Aequationes Mathematicae*, 47 (1994), 293.
- [24] Yu.L. Shmul’yan, Two-sided division in a ring of operators, *Math. Notes* 1, (1967), 400–403.

- [25] B. Simon, Lower semi-continuity of positive quadratic forms, Proc. Roy. Soc. Edinburgh, 79 (1977), 267-273.
- [26] B. Simon, A canonical decomposition for quadratic forms with applications to monotone convergence theorems, J. Functional Analysis, 28 (1978), 377-385.
- [27] J. Stochel and F.H. Szafraniec, *Unbounded operators and subnormality*, Manuscript, 2001.
- [28] M.H. Stone, On unbounded operators in Hilbert space, J. Indian Math. Soc., 15 (1951), 155-192.
- [29] W. Szymański, Decomposition of operator-valued functions in Hilbert spaces, Studia Math., 50 (1974), 265-280.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VAASA, P.O. Box 700, 65101 VAASA, FINLAND

E-mail address: `sha@uwasa.fi`

DEPARTMENT OF APPLIED ANALYSIS, EÖTVÖS LORÁND UNIVERSITY, PÁZMÁNY PÉTER SÉTÁNY 1/C, 1117 BUDAPEST, HUNGARY

E-mail address: `sebesty@cs.elte.hu`

DEPARTMENT OF MATHEMATICS AND COMPUTING SCIENCE, UNIVERSITY OF GRONINGEN, P.O. BOX 800, 9700 AV GRONINGEN, NEDERLAND

E-mail address: `desnoo@math.rug.nl`

INSTITUTE OF MATHEMATICS, UNIERSYTET JAGIELLOŃSKI, UL. REYMONTA 4, 30059 KRAKÓW, POLAND

E-mail address: `fhszafra@im.uj.edu.pl`